

NIH Resources

The NINDS also strongly encourages researchers to use these NIH developed materials for NINDS-sponsored research, when appropriate. Utilization of these resources will enable greater consistency for NINDS-sponsored research studies. These tools are free of charge.

- National Institutes of Health (NIH) Toolbox
- Quality of Life in Neurological Disorders (Neuro-QOL)
- Patient-Reported Outcomes Measurement Information System (PROMIS)

Suicidal Ideation

Investigators should review the FDA's "Guidance for Industry: Suicidal Ideation and Behavior: Prospective Assessment of Occurrence in Clinical Trials" for the most up-to-date information about suicidal ideation and behavior. One scale that FDA suggests is the Columbia Suicide Severity Rating Scale (C-SSRS) (available at http://www.cssrs.columbia.edu

Disease/Domain	Recommendations
Facioscapulohumeral muscular dystrophy	These instruments and elements are recommended for use in FSHD studies:
	Core elements:
	See Start-Up Resources Listing document (FSHD Start-Up Resource Listing)
	Supplemental – Highly Recommended instruments: 6 Minute Walk Test, Grip Strength ⁱ , Manual Muscle Testing-Using the Medical Research Council Muscle Grading Scale, Maximum Voluntary Isometric Contraction Testing (MVICT)
Participant / Subject	
Participant / Subject	Core: General Core, Demographics ⁱⁱ
Characteristics;	Core: General Core, Demographics ⁱⁱ Supplemental – Highly Recommended: None
Characteristics;	Supplemental – Highly Recommended: None
Characteristics;	Supplemental – Highly Recommended: None Supplemental: Demographics
Characteristics; Demographics Participant / Subject Characteristics; Social	Supplemental – Highly Recommended: None Supplemental: Demographics Exploratory: None
Characteristics; Demographics Participant / Subject	Supplemental – Highly Recommended: None Supplemental: Demographics Exploratory: None Core: None



Disease/Domain	Recommendations
Participant/Subject History and Family History; Epidemiology/ Environmental History	Core: None Supplemental – Highly Recommended: None Supplemental: None Exploratory: World Health Organization Disability Assessment Schedule (WHODAS)
Participant/Subject History and Family History; General Health History	Core: None Supplemental – Highly Recommended: None Supplemental: Family History, Medical History, Pregnancy Outcome Exploratory: None
Assessments and Examinations; Imaging Diagnostics	Core: None Supplemental – Highly Recommended: None Supplemental: Dual X-Ray Absorptiometry (DXA), Magnetic Resonance Imaging (MRI) Exploratory: None
Assessments and Examinations; Laboratory Tests and Biospecimens/ Biomarkers	Core: None Supplemental – Highly Recommended: Laboratory Tests and Tracking Supplemental: Mutation Analysis Exploratory: None
Assessments and Examinations; Non-Imaging Diagnostics	Core: None Supplemental – Highly Recommended: Electrocardiogram (ECG) Supplemental: Electrocardiogram (ECG) Exploratory: None
Assessments and Examinations; Physical/Neurological Examination	Core: None Supplemental – Highly Recommended: None Supplemental: Polysomnography Exploratory: Electrical Impedance Myography (EIM)
Treatment/ Intervention Data; Drugs	Core: None Supplemental – Highly Recommended: Prior and Concomitant Medications Supplemental: None Exploratory: None



Disease/Domain	Recommendations
Treatment/ Intervention Data; Therapies	Core: None
	Supplemental – Highly Recommended: None
	Supplemental: Respiratory Interventions
	Exploratory: None
Outcomes and Endpoints; Functional Status	Core: None
	Supplemental – Highly Recommended: 6 Minute Walk Test
	Supplemental: Goniometry, Stair Climb, Timed Up and Go (TUG)
	Exploratory: 2 Minute Walk Test
Outcomes and Endpoints; Muscle Strength Testing	Core: None
	Supplemental – Highly Recommended: Grip Strength, Manual Muscle Testing- Using the Medical Research Council Muscle Grading Scale, Maximum Voluntary Isometric Contraction Testing (MVICT)
	Supplemental: Hand Held Dynamometry, Pinch Strength
	Exploratory: None
Outcomes and Endpoints; Pain	Core: None
	Supplemental – Highly Recommended: None
	Supplemental: Brief Pain Inventory (Short Form), McGill Pain Questionnaire
	Exploratory: None
Outcomes and Endpoints; Pulmonary Function Testing/Respiratory Status	Core: None
	Supplemental – Highly Recommended: Pulmonary Function Testingiv
	Supplemental: Measures of Gas Exchange
	Exploratory: None
Outcomes and Endpoints; Quality of Life	Core: None
	Supplemental – Highly Recommended: None
	Supplemental: Individualized Neuromuscular Quality of Life Questionnaire (INQoL), Short Form 36-Item Health Survey-36 (SF-36)
	Exploratory: Quality of Life in Neurological Disorders (NeuroQoL)

ⁱ Supplemental – Highly Recommended for Facioscapulohumeral Muscular Dystrophy (FSHD) studies involving muscle strength testing.

FSHD CDE Highlight Summary Last Modified 2/13/14

ii Contains Core CDEs



[&]quot;To be completed by female participants/subjects only.

The most useful measure of pulmonary function for FSHD is sitting and supine Forced Vital Capacity. Pulmonary Function Testing (PFT) is likely to have the greatest yield in clinical studies in which subjects are non-ambulatory and have severe weakness or early ("infantile") onset of disease. PFT should be performed with a face mask in FSHD due to the inability to form a tight seal with a mouthpiece.